

# Localization by frustration: From magnons to electrons

J. Richter<sup>a</sup>, O. Derzhko<sup>b</sup> and A.Honecker<sup>c</sup>

<sup>a</sup>ITP, Universität Magdeburg, P.O. Box 4120, D-39016 Magdeburg, Germany

<sup>b</sup>Institute for Condensed Matter Physics, L'viv-11, 79011, Ukraine

<sup>c</sup>Institut für Theoretische Physik, Universität Göttingen, D-37077 Göttingen, Germany

For antiferromagnetic spin systems as well as for Hubbard electrons on various frustrated lattices a class of exact eigenstates can be constructed. Such eigenstates can be found, e.g., for the 1D sawtooth chain, the 2D kagomé and checkerboard, and the 3D pyrochlore lattices. The exact many-particle eigenstates consist of independent magnons/electrons localized on finite areas of the lattice. Important structural elements of the relevant systems are triangles being attached to polygons or lines. Then the magnons/electrons can be trapped on these polygons/lines. For electrons the scenario of localized eigenstates is related to the so-called flat-band ferromagnetism.

The correlated systems having localized eigenstates exhibit a highly degenerate ground-state manifold at the saturation field  $h_{sat}$  (at a characteristic value of the chemical potential  $\mu_0$ ) for magnons (electrons). The degeneracy grows exponentially with the system size and leads to a finite residual entropy. By mapping the localized magnon (electron) degrees of freedom onto a hard-core lattice gas one may find explicit analytical expressions for the low-temperature thermodynamics in the vicinity of  $h_{sat}$  ( $\mu_0$ ). Though the scenario of localized eigenstates is similar for spin and electron systems, the different statistics of spins and electrons leads to different construction rules for the localized eigenstates and, as a result, to a different hard-core lattice gas description.

9.7 cm

13.4 cm

## Subject category :

2. Quantum and Classical Spin Systems

## Presentation mode :

oral

## Corresponding author :

J. Richter

## Address for correspondence :

Institut für Theoretische Physik, Universität Magdeburg,  
P.O. Box 4120, D-39016 Magdeburg, Germany

## Email address :

Johannes.Richter@Physik.Uni-Magdeburg.DE